

Hidden reservoirs of pathogens in dental settings

Silpi Chatterjee^{1,*}, Sonal Saigal², Ankur Bhargava³, Daya Shankar⁴, Asim Mustafa Khan⁵, & Safiya Fatima Khan⁶

¹Department of Public Health Dentistry, Hazaribag college of Dental Sciences and Hospital, Jharkhand, India; ²Department of Oral Pathology and microbiology and forensic odontology, Dental Institute, RIMS, Ranchi, India; ³Department of Oral Pathology and microbiology and forensic odontology, Hazaribag college of Dental Sciences and Hospital, Jharkhand, India; ⁴Department of rosthodontics and Crwon & Bridge. Hazaribag college of Dental Sciences and Hospital, Jharkhand, India; ⁵Department of Biomedical Dental Sciences Imam Abdulrahman Bin Faisal University; ⁶Department of Periodontology Faculty of Dental Sciences, Ramaiah University of Applied Sciences; Dr Silpi Chatterjee – E-mail: dr.silpi510@gmail.com; Corresponding author*

Received December 5, 2020; Revised December 31, 2020; Accepted January 2, 2020, Published January 31, 2021

DOI: 10.6026/97320630017073

Declaration on Publication Ethics:

The author's state that they adhere with COPE guidelines on publishing ethics as described elsewhere at <https://publicationethics.org/>. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues connecting to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher in regard to this article.

Author responsibility:

The authors are responsible for the content of this article. The editorial and the publisher have taken reasonable steps to check the content of the article in accordance to publishing ethics with adequate peer reviews deposited at PUBLONS.

Declaration on official E-mail:

The corresponding author declares that official e-mail from their institution is not available for all authors

Abstract:

Nosocomial infections are a major concern to both clinicians and health care seekers. Investigations have suggested that laptops & mobile phones may contribute to cross-contamination and can serve as vehicles for infection transmission. Therefore, it is of interest to document the data on hidden reservoirs such as mobile phones and laptops of pathogens in dental settings at the Hazaribag College of dental sciences and Hospital, Jharkhand. The samples were collected from 25 laptops and 25 mobile phones from dentists working in a dental college in Hazaribag city. The samples were collected aseptically using sterile cotton swabs dipped in sterile saline by rotating the swabs on the keyboard surfaces of laptops and mobile phones, inoculated into Brain Heart Infusion broth, vortexed for 1 minute in Fischer Vortex Genie 2 on highest setting & streaked immediately on 5% sheep blood agar plates and were incubated at 37°C for 24 hours aerobically. The isolates were identified based on the colony morphology, colony characteristics and biochemical reactions. The bacterial species isolated were *Staphylococcus aureus*, Coagulase negative *Staphylococcus*, *Bacillus species*, *Enterococci*, *Micrococci*, and *Pseudomonas* etc. Predominant species isolated was *Staphylococcus aureus* and least was *Micrococci*. Higher percentage of organisms was found at the Department of Periodontics, Endodontics and least was found in Department of Public Health Dentistry. The percentage and type of organism isolated from keyboards of laptops and mobile phones were similar. Thus, laptops and mobile phones act as vehicles for transfer of potential pathogens associated with dental hospitals. Disinfecting the hands prior to examination of patients and disinfection of laptops and mobiles with alcohol wipes should be done to prevent nosocomial infections.

Key words: Disinfection; infection; microbial contamination; pathogenic organism.

Background:

Healthcare-associated infections are an important cause of morbidity and mortality in hospitals. Each year more than 2 million patients acquire healthcare-associated infections, resulting in 90,000 deaths and healthcare costs that are estimated to exceed \$5 billion. Health care-associated infection (HCAI), also referred to as “nosocomial” or “hospital” infection, is defined as: “An infection occurring in a patient during the process of care in a health-care facility which was not present or incubating at the time of admission” [1]. Some studies have demonstrated that the mean rate of compliance with the Centers for Disease Control and Prevention guidelines on hand hygiene is approximately 40% among healthcare workers [2], which is a likely explanation for the frequent contamination of computer keyboards and mobile phones. With the advent of technology, mobile phones, laptops used by health care professionals are on the rise especially in the clinical set ups. The laptops and mobile phones of health care workers harbor many harmful pathogens which serve as a reservoir for nosocomial infections and may contribute to cross – contamination, which serve as vehicles for infection transmission [3,6]. Studies have revealed that mobile phones and laptops have a great potential for dissemination of disease and the incidence of such cross contamination diseases to be 4.8% in U S A, 7.1% in European countries, 10-30% in India and 17.1 % in Iran [7,10]. Some investigators have suggested that computer keyboards may contribute to cross-transmission because of acquisition of transient hand carriage by healthcare personnel during contact with the contaminated computer keyboard surface [11,12]. Technical support systems have acted as a boon for health care providers in the past few decades. The burden of data recording, data maintenance & analysis of data have become very easy with the introduction of multiple softwares in health care sector. The usage of these has been very simple & can be operated through laptops & mobile phones. This in turns acts as reservoir for health care associated infections. Since laptops and mobile phones have become an essential means of communication, their usage in clinical set up is unavoidable [13]. As mobile phones act as perfect habitat for microbes to breed, especially in high temperature and humid conditions, Health care workers (HCWs’) mobile phones may serve as reservoirs of microorganisms that could be easily transmitted from the mobile phones to the HCWs’ hands and therefore facilitate the transmission of bacterial isolates from one patient to another in different hospital wards [14]. Dental clinics are commonplace for the bacterial aerosols generated by high-speed dental hand pieces with water supplies, which has the capacity to settle over long distance. Aerosols and spatter produced during many dental procedures are a potential source of transmission of various diseases [15,17]. The use of laptops, desktops, and mobile

phones has become an integral part of dental practice. Therefore, it is of interest to document the data on hidden reservoirs such as mobile phones and laptops of pathogens in dental settings at the Hazaribag College of dental sciences and Hospital, Jharkhand.

Methodology:**Study location:**

A cross sectional study was done to assess the microbial contamination of laptops and mobile phones used by dentists in clinical settings of a dental college in Hazaribag city, Jharkhand, India.

Study Period:

The duration of the study was for a period of 3 months from January 1st to March 31st 2020.

Ethical Clearance:

Ethical clearance was obtained from the Institutional Ethical Committee before the start of the study. Necessary permission was obtained from the institution prior to the study. Informed consent was obtained from the dentists before the start of the study.

Study Criteria:

Inclusion criteria were the laptops and mobile phones, which were in use for a minimum period of one year near clinical settings, were taken for the study.

Consent:

Dentists who did not give consent to participate were excluded from the study.

Data size:

A pilot study was conducted by collecting the samples **from 5 participants**.

Model data:

A sample of 25 laptops and 25 mobile phones, which satisfied eligibility criteria, were considered for the study. The laptops and mobile phones were randomly selected using a simple random sampling technique.

Microbial analysis:

The samples were collected aseptically using sterile cotton swabs dipped in sterile saline by rotating the swabs on the keys of laptops and mobile phones during operating hours using a method in which the investigator had received training in advance. The swabs were then transported immediately to the laboratory for inoculation. The samples were then inoculated into Brain Heart

Infusion (BHI) broth. The sample was vortexed for 1 minute in Fischer vortex genie 2 on highest setting. The samples were then streaked immediately on 5% sheep blood agar plates and were incubated at 37°C for 24 hours aerobically. The organisms isolated were stained and identified based on their morphology (shape, arrangement of the organisms), colony characteristics (size, shape of the colony, opacity, pigmentation, haemolysis, elevation etc.) and biochemical reactions (catalase test, coagulase test, sugar fermentation, heat test, citrate utilization test, urease test, triple

sugar iron test, oxidase, mannitol motility test etc). The colonies were counted and colony-forming unit was estimated.

Statistical analysis:

The data analysis was done using the statistical software SPSS version 23. Descriptive statistics was done for the colony forming units and microbial organisms present on laptops and mobile phones of various departments. Pearson correlation was computed for comparing the microbial contamination of laptops & mobile phones with respect to various departments.

Table 1: Distribution of microbial contamination of laptops and mobile phones according to speciality.

Department	Laptops CFU-N (%)	Mobiles CFU-N (%)	Total CFU per dept	% (Total CFU per dept/ Total CFU of all departments)	Total CFU of all departments
Public Health Dentistry	3900(6.19)	1600(4.74)	5500	5.68	
Pedodontics	8100(12.85)	1300(3.85)	9400	9.71	
Prosthodontics	8800(13.96)	2550(7.55)	11350	11.72	
Oral surgery	9200(14.6)	6250(18.51)	15450	15.95	
Oral medicine	3300(5.23)	1000(2.96)	4300	4.44	
Oral pathology	3500(5.55)	800(2.37)	4300	4.44	
Endodontics	10500(16.66)	3600(10.66)	14100	14.55	
Periodontics	6400(10.15)	8600(25.48)	15000	15.5	
Orthodontics	9300(14.76)	8200(24.29)	17500	18.05	96900

Table 2. Distribution of microbial contamination of mobiles and laptops among specialities

Organism	Public Health	Pedodontics	Prosthodontics	Oral surgery	Oral medicine	Oral pathology	Endodontics	Periodontics	Orthodontics	
Staphylococcus aureus	Mobile	0	4.17	10	16.6	0	2.5	16.7	29.2	20.8
	Laptop	0	1.25	12.5	18.75	10.63	10	22.5	25	21.87
Coagulase -ve staphylococcus	Mobile	29.63	9.88	17.28	8.64	12.3	2.4	9.88	0	9.87
	Laptop	2.56	5.12	16.67	3.84	11.54	10.3	25.6	5.13	19.23
Micrococci	Mobile	12	8	8	4	0	0	12	28	28
	Laptop	0	0	31.03	13.79	3.44	0	6.9	17.2	27.58
Acinetobacter	Mobile	0	0	12.5	0	0	0	50	0	0
	Laptop	33.87	0	19.35	27.42	0	0	1.61	11.3	6.45
Diphtheroids	Mobile	0	0	33.33	0	0	0	66.7	0	0
	Laptop	0	22.73	15.91	13.64	4.54	0	22.7	11.4	9.09
B.Anthraxis	Mobile	0	0	0	0	100	0	0	0	0
	Laptop	0	6.66	13.33	10	0	0	23.3	33.3	13.33
B.subtilis	Mobile	3.85	3.85	3.84	0	0	15.4	3.85	65.4	3.84

	Laptop	16.12	41.94	1.07	8.6	0	10.8	3.23	16.1	2.15
Enterococci	Mobile	0	0	0	0	9.5	0	0	66.7	23.81
	Laptop	0	0	7.89	15.79	0	0	18.4	7.89	13.15
Pseudomonas	Mobile	0	20	70	60	20	0	0	0	0
	Laptop	0	0	11.76	23.53	0	0	5.88	13.5	35.29

Results:

The organisms isolated were *Staphylococcus aureus*, Coagulase negative Staphylococci, *Micrococci*, *Enterococci*, *Diphtheroids*, *Bacillus anthracis*, *Bacillus subtilis*, *Acinetobacter species* and *Pseudomonas species*. Out of 25 laptops, 16.66% of laptops from the Department of Endodontics, 14.76% from Department of Orthodontics, 14.6% from Department of Oral surgery were contaminated. Among mobiles, 25.48% of mobiles from the Department of Periodontics, 24.29% from Orthodontics, 18.51% from Oral surgery were contaminated (**Table 1**). *Staphylococcus aureus* was present in 29.2% of mobiles from Periodontics, 20.8% from Orthodontics and 16.7% from Endodontics and 16.6% from Oral surgery. Coagulase negative *Staphylococcus* was present in 29.63% of mobiles from Public Health Dentistry, 17.28% from Prosthodontics and 12.3% from Oral medicine (**Table 2**). *Staphylococcus aureus* was present in 22.5% of laptops from Endodontics followed by 21.87% from Orthodontics, 18.75% from Oral surgery and 12.5% from Prosthodontics. Coagulase negative *Staphylococcus* was present in 25.6% from Endodontics, 19.23% from Orthodontics, 16.67% from Prosthodontics and 11.54% from Oral medicine (**Table 2**). *Staphylococcus aureus* was present in all the laptops (88.89%) and (77.78%) of mobiles (**Table 3**). Statistically significant and positive correlation was obtained for department of Prosthodontics ($r=0.809$), Oral pathology ($r=0.894$) and Endodontics ($r=0.860$) (**Table 4**).

Table 3. Distribution of microorganisms in laptops and mobiles

Microorganism	% of microbial contamination of laptops	% of microbial contamination of mobiles
<i>Staphylococcus aureus</i>	88.89	77.78
Coagulase negative staphylococcus	100	88.89
Micrococci	66.67	77.78
Enterococci	66.67	33.34
Diphtheroids	77.78	22.23
<i>B. anthracis</i>	66.67	11.12
<i>B. subtilis</i>	88.89	77.78
<i>Acinetobacter species</i>	62.5	22.23
<i>Pseudomonas species</i>	66.67	33.34

Table 4: Correlation between laptops and mobile phones among various specialities

*Statistically significant at $p < 0.05$

Department (Laptop * Mobile)	r value	p value
Public Health Dentistry	-0.164	0.63

Pedodontics	-0.108	0.751
Prosthodontics	0.809*	0.003*
Oral surgery	-0.96	0.779
Oral medicine	0.395	0.229
Oral pathology	0.894*	0.000**
Endodontics	0.860*	0.001*
Periodontics	0.323	0.333

Discussion:

This study shows that a proportion of around two third of all the laptops and mobile phones near clinical setup and almost half of those sampled immediately after use were contaminated with microorganisms, which can lead to nosocomial infections. The microbial contamination was more for the departments of Orthodontics (18.08%) followed by Oral surgery (15.96%) and least was from Endodontics (14.57%). The use of mobile phones and laptops by the dental faculty and postgraduate students involved in direct patient care not only demonstrated a high contamination rate with bacteria but were contaminated with nosocomial pathogens. The organisms isolated were *Staphylococcus aureus*, *Micrococci*, *Acinetobacter*, *Bacillus species*, *Enterococci* and *Pseudomonas*. Among these *Staphylococcus aureus* and *Acinetobacter* are resistant to drying and can survive for weeks in a dry environment and is capable of multiplying rapidly. *Acinetobacter* was identified based on Gram stain, oxidase and motility tests. The microbial contamination in the present study among laptops was *Staphylococcus aureus* 25% from department of Periodontics followed by 22.5% from the department of Endodontics, coagulase negative *Staphylococcus* was 25.6% from the department of Endodontics which is contradictory to study on laptops which showed 88% of contamination with coagulase negative staphylococcus and 12% of contamination with *staphylococcus aureus* [17]. The overall rate of contamination of laptops with potentially pathogenic organisms like *Acinetobacter* was 62.50% which is similar to a study by William A et al [18] where multidrug resistant *Acinetobacter baumannii* was found on the hands, cell phones of health care workers and patients admitted to the ICU (60%) and contradictory to a study by Sweta Singh et al. [19] on cell phones which showed lower rates of contamination ranging from 7-14.3%. The higher rates of contamination of laptops and mobile phones among departments in this present study might be due to the influence of various factors like lack of hand washing after examination or treatments, disinfection practices followed in

the hospital, frequency of use of gadgets and the frequency of disinfection of laptops and mobile phones. This study showed that 88% of laptops and 98% of the mobile phones were contaminated with more than one pathogenic organism which is similar to a study done by Brady *et al.* [4] showed that 89.7% of mobile phones were contaminated. The most dominant organism isolated was *Staphylococcus aureus*. Jesle *et al.* [20] found that rate of bacterial contamination of hospital care workers (HCW's) was 95% while that of mobile phones was 90% which is similar to a study by Sweta Singh *et al.* [19] who reported that out of 50 mobile phones cultured, 98% were positive. The present study is contradictory to a study by Harish Trivedi *et al.* [21] where 58.66% of hand samples and 46.66% of mobile phones were contaminated by bacteria. Ulger *et al.* [22] showed that 94.5% of phones showed evidence of bacterial contamination. They found that 49% of phones had one bacterial species, 34% had two different species and 11.5% had two or more different species which is contradictory to the present study were 20% had single species (n=3), 45% had two species (n=15) and 35% had more than two types of species (n=7). However, lower rates were observed by Ramesh *et al.* [23] in which, 45% of mobile phones were contaminated. A study by Lu *et al.* [24] revealed a 17.4% contamination rate of computer devices by *Staphylococcus aureus*, *Acinetobacter species* or *Pseudomonas species* and contradictory to a study by William *et al.* [18] who studied the degree of microbial contamination of computers, the efficacy of different disinfectants, and the cosmetic and function effects of these disinfectants on computer keyboards. Potential pathogenic microorganisms were cultured from more than 50 percent of the computers. 10.15% of laptops from Department of Periodontics were contaminated followed by 13.96% from Prosthodontics, 16.6% from Endodontics. In case of mobile phones, 25.48% from Periodontics, 10.66% from Endodontics were contaminated, 7.55% from Prosthodontics which is contrast to a study by Sham.S.Bhat *et al.*²⁵ where 4 % from Prosthodontics, 5% of mobiles from Orthodontics showed pathogenic organisms. Hence, in a country like India, mobile phones and laptops of HCWs plays an important role in transmission of infection to patients, which can increase the burden of health care. Simple measures such as increasing hand hygiene and regular decontamination of mobile phones with alcohol disinfectant wipes may reduce the risk of cross contamination caused by these devices. One study reported the use of 70 % isopropyl alcohol as an effective disinfectant¹⁵. Another study reported that restricted use of mobile phones during working hours along with proper hand hygiene practices enabled mobile phones to remain free of contamination¹⁶. The findings of the present study are alarming which shows that dentists are lacking the awareness of the safety measures when a significant number of them neither clean their hands before and after seeing a patient nor disinfect

their laptops and mobile phones after using in the hospital setup. Hand washing is the simplest and most economical measure that can prevent the transfer of harmful pathogens. There are no rules restricting dentists to use laptops and mobile phones into a sterile clinical setup in India. There are also no cleaning guidelines for laptops and mobile phones of health care workers. The design of this study being a cross-sectional one does not permit causal inference between microorganisms present in laptops and mobiles. Further studies for the assessment of microbial contamination among dental specialties and methods of decontamination of laptops and mobile phones should be formulated.

Conclusion:

Laptops and mobile phones are reservoir of microorganisms associated with healthcare associated infections (HAI). Data shows that 88% of laptops and 98% of the mobile phones were contaminated. It appears that routine disinfection of mobile phones and laptops will be effective in reducing microbial contamination.

References:

- [1] The Burden of Health Care-Associated Infection Worldwide. A Summary World Health Organization.
- [2] Harrel SK and Molinari J, *J Am Dent Assoc.* 2004 **135**:429.
- [3] Pina-Vaz *et al. Rev Clin Pesq Odontol.* 2008 **4**:77.
- [4] Brady RR *et al. J Hosp Infect.* 2007 **66**:397.
- [5] M.Shakeel *et al. International Journal of Public Health Dentistry* 2011 **2**:4.
- [6] Edwards A *et al. J Hosp Infect* 2002 **52**:314.
- [7] Palmer SR and Bray SL. *Aust J Educ Technol* 2001 **17**:313.
- [8] Bures S *et al. Am J Infect Control* 2000 **28**:465.
- [9] Prathibha P *et al. RRJDS* 2013 **1**:1.
- [10] Mir Mehdi F Study of the nosocomial infections in Tabriz Imam Hospital, the 11th congress of infectious and tropical diseases in Iran, 2002 46.
- [11] Oguz Karabay *J Infect Developing Countries* 2007 **1**:72.
- [12] Maureen Schultz Janet Gill *et al. Infection control and hospital epidemiology* 2003 **24**:302.
- [13] Harrel SK and Molinari J *J Am Dent Assoc.* 2004 **135**:429.
- [14] Rawia Ibrahim Badr *et al. Int J Infect Control* 2012 **8**:1.
- [15] Leggat PA *et al. Int Dent J.* 2001 **51**:39.
- [16] Leggat PA and Kedjarune U *Ind Health.* 2007 **45**:611.
- [17] Szymanska J *Ann Agric Environ Med.* 2007 **14**:203.
- [18] William A *et al. Infection control and hospital epidemiology* 2006 **27**:372.
- [19] Sweta Singh *et al. Journal of Dental Education* 2010 **74**:1153.
- [20] Jesle H-C *et al. Anaesthesia* 2007 **62**:904.
- [21] Dr Harish R *et al. NJIRM* 2011 **2**:61.
- [22] Ulger F *et al. Ann Clin Microbiol Antimicrob* 2009 **8**:7.

[23] Ramesh J *et al.* *Journal of Hospital Infection* 2008 **70**:160.
[24] Lu PL *et al.* *BMC infect Dis* 2009 **9**:164.

[25] Sham S *et al.* *Online Journal of Health and Allied Sciences* 2011 **10**:1.

Edited by P Kanguane


Citation: Chatterjee *et al.* *Bioinformation* 17(1): 73-79 (2021)

License statement: This is an Open Access article which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. This is distributed under the terms of the Creative Commons Attribution License

Articles published in BIOINFORMATION are open for relevant post publication comments and criticisms, which will be published immediately linking to the original article for FREE of cost without open access charges. Comments should be concise, coherent and critical in less than 1000 words.

BIOINFORMATION

Discovery at the interface of physical and biological sciences



since 2005

BIOINFORMATION

Discovery at the interface of physical and biological sciences

indexed in

