

Molecular docking of alkaloid compounds with the matrix metalloproteinase 2

JH Shazia Fathima¹, Jayaraman Selvaraj J^{2*}, Venkatalalam Sivabalan³, Umapathy Vidhya Rekha⁴, Rajagopal Ponnulakshmi⁵, Veeraraghavan Vishnupriya², Malathi Kullappan⁶, Radhika nalinakumari Sreekandan⁷, Surapaneni Krishna Mohan⁸

¹Department of Oral and Maxillofacial Pathology, Ragas Dental College and Hospitals, Chennai, India; ²Department of Biochemistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai - 600 077, India; ³Department of Biochemistry, KSR Institute of Dental Sciences and Research, Thiruchengodu-637215, India; Department of Public Health Dentistry, Sree Balaji Dental College and Hospital, Pallikaranai, Chennai-600 100, India; ⁵Central Research Laboratory, Meenakshi Academy of Higher Education and Research (Deemed to be University), Chennai-600 078, India; ⁶Department of Research, Panimalar Medical College Hospital & Research Institute, Varadharajapuram, Poonamallee, Chennai - 600 123, India; ⁷Department of Clinical Skills & Simulation, Panimalar Medical College Hospital & Research Institute, Varadharajapuram, Poonamallee, Chennai - 600 123, India; ⁸Department of Biochemistry and Department of Clinical Skills & Simulation, Department of Research, Panimalar Medical College Hospital & Research Institute, Varadharajapuram, Poonamallee, Chennai - 600 123; ⁹DBT-BIF Centre, PG & Research Department of Biotechnology & Bioinformatics, Holy Cross College (Autonomous), Trichy, Tamilnadu, India; *Corresponding Author: Dr. Jayaraman Selvaraj - E-mail: jselvaendo@gmail.com

Author contacts:

Shazia Fathima JH-shaziafathimarizwan@gmail.com; Selvaraj Jayaraman- jselvaendo@gmail.com; Venkatalalam Sivabalan - biosivabalan@gmail.com; Umapathy Vidhya Rekha- drvidhyarekha@gmail.com; Rajagopal Ponnulakshmi-ramgslaks@gmail.com; Veeraraghavan Vishnupriya-drvisnupriyav@gmail.com; Malathi Kullappan-malak.hari@gmail.com; Radhika Nalinakumari Sreekandan-niharakrishna21@gmail.com; Surapaneni Krishna Mohan -krishnamohan.surapaneni@gmail.com;

Received December 29, 2020; Revised December 31, 2020; Accepted January 26, 2020, Published January 31, 2021

DOI: 10.6026/97320630017206

Declaration on Publication Ethics:

The author's state that they adhere with COPE guidelines on publishing ethics as described elsewhere at <https://publicationethics.org/>. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues connecting to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher in regard to this article.

Author responsibility:

The authors are responsible for the content of this article. The editorial and the publisher have taken reasonable steps to check the content of the article in accordance to publishing ethics with adequate peer reviews deposited at PUBLONS.

Declaration on official E-mail:

The corresponding author declares that official e-mail from their institution is not available for all authors

Abstract:

Matrix metalloproteinase protein-2 (MMP-2) is linked to the human oral squamous cell carcinoma. Therefore, it is of interest to design new inhibitors for MMP-2 to combat the disease. Thus, we document the molecular docking features of Aristolochic acid, Cryptopleurine, Epipodophyllotoxin, and Fagaronine with MMP-2 for further consideration.

Key words: MMP-2, OSCC, alkaloids, molecular docking

Background:

Matrix metalloproteinase protein-2 (MMP-2) is linked to the human oral squamous cell carcinoma [1-6]. Data shows the role of MMP-2 as a cancer prognostic marker [7] for prognosis [8]. Therefore, it is of interest to design new inhibitors for MMP-2 to combat the disease.

Table 1: List of selected alkaloids compounds

S.No	Compound Name	Source of plant
1	Aristolochic acid	Aristolochia indica L.
2	Camptothecin	Camptotheca acuminata Decne
3	Colchicine	Colchicum speciosum Steven
4	Cryptopleurine	Crinum macrantherium Engl
5	Epipodophyllotoxin	Podophyllum species L.
6	Demecolcine	Colchicum speciosum Steven
7	Fagaronine	Fagara zanthoxyloides Lam
8	Oxyacanthine	Berberis asiatica Roxb.
8	Thalicarpine	Thalictrum dasycarpum Fisch

Molecular Docking:

The PyRx Version 0.8 [11,12] was used for molecular docking analysis. The results were visualized using Pymol.

Results and Discussion:

Alkaloids seem to be significant chemical compounds for drug development. Alkaloids derived from natural herbs exhibit antiproliferation and antimetastasis effects on different forms of cancers. A significant component of the chemotherapy arsenal is its use for cancer care. The role of alkaloids appears to be unique in the cell cycle. Therefore, certain alkaloid compounds from various medicinal plants are used to assess their effectiveness against MMP-2 in OSCC. The top four best-docked compounds with a stronger affinity for the MMP-2 receptor were chosen, according to binding mode and molecular interaction analysis in the MMP-2 binding cavity (Table 2). Molecular interaction and docking

Table 2: Molecular interaction obtained from PyRx

S.No	Compound Name	Binding Energy	H-bond Interaction	Distance Å
1	Aristolochic acid	-7.1	PHE-87	2.9
			HIS-124	2.4
			GLU-129	2.9
			HIS-130	2.5
2	Cryptopleurine	-6.4	ALA-88	2.5
3	Epipodophyllotoxin	-6.2	ALA-88	2.4
4	Fagaronine	-6	HIS-124	2.1

Materials and Methods:**Ligand Preparation:**

Ligands are downloaded from the Pubchem database (Table 1) [9] and converted into 3D data using Pymol. The data was stored for AutoDock vina-PyRx in pdb file format. Ligand data were then saved for PyRxx in the PDBQT file format.

Protein Preparation:

The protein PDB ID 1HOV for MMP-2 was downloaded from the RCSB protein database PDB for this study and prepared according to a standard procedure [10].

measurements indicate that the lead compound Aristolochic acid offers -7.01 kcal/mol binding energy and has been shown to be more effective as an inhibitor of MMP-2 against MMP-2. Hydrogen bonding is an interaction reaction whereby the donors of hydrogen bonds and free protein and ligand acceptors sever their hydrogen bonds with water and in the protein-ligand complex form new ones. In addition to filtering unrealistic poses in docking, hydrogen-bonding formation could also be used to increase the precision of binding energy measurement. The Aristolochic acid molecular interaction analysis offers insights into their binding mode with the MMP-2 receptor and the MMP-2 amino acids that evaluate the efficacy of the docking compound. Four hydrogen bonds formed by PHE-87, HIS-124, GLU-129, and HIS-130 with bond lengths of 2.9\AA , 2.4\AA , 2.9\AA , 2.5\AA have been formed by aristolochic acid (Figure 1a). There were less than three H-bonds, suggesting the existence of favourable associations between the

ligand and the receptor. The association of cryptopleurin with the protein MMP-2 was shown in **Figure 1b**. The amino acid residue ALA-88 formed the hydrogen bond interaction with MMP-2 at a distance of 2.5 Å. It has been calculated that cryptopleurine docking energy with MMP-2 is 6.4 Kcal / Mol. **Figure 1c** shows the docking conformation of Epipodophyllotoxin with MMP-2. The amino acids related to the interaction of the hydrogen bond with MMP-2 reacting with Epipodophyllotoxin were determined to be ALA-88

and the bonding wavelength of hydrogen is 2.4Å. The docking energy was computed and identified to be 6.2 Kcal/Mol for Epipodophyllotoxin with MMP-2. The docking conformation of Fagaronine with MMP-2 is shown in **Figure 1d**. Bonded hydrogen bond with residues HIS-124 of MMP-2 with a distance of 2.1Å is reinforced by Fagaronine with MMP-2 complex. It was observed that the binding energy for Fagaronine to MMP-2 was-6.0 Kcal / Mol.

Figure needed in single image file

Figure 1: Molecular docking interaction of MMP-2 with (a) Aristolochic acid; (b) Cryptopleurine; (c) Epipodophyllotoxin; and (d) Fagaronine

Conclusion:

We document the molecular docking features of Aristolochic acid, Cryptopleurine, Epipodophyllotoxin, and Fagaronine with MMP-2 for consideration in the context of OSCC.

References:

- [1] Markopoulos AK, *Open Dent J.* 2012 **126**. [PMID: 22930665].
- [2] Scully C *et al. Oral Oncol.* 2009 **45**:301. [PMID: 19249237].
- [3] Gillison ML. *Head Neck.* 2007 **29**:779. [PMID: 17230556]
- [4] Petti, S, *Oral Oncology.* 2009 **45**:340. [PMID: 18674956].
- [5] Rosebush MS *et al. J Tenn Dent Assoc.* 2011 **91**:24. [PMID: 21748976].
- [6] Roomi MW *et al. Oncol Rep.* 2009 **21**:1323. [PMID: 19360311].
- [7] Chen Y *et al. Carcinogenesis.* 2014 **35**:442. [PMID: 24072772].
- [8] Wen X *et al. Tumour Biol.* 2014 **35**:845. [PMID: 24037915]
- [9] Kim S *et al. Nucleic Acids Res.* 2019 **47**:D1102. [PMCID: PMC6324075]
- [10] Bernstein FC *et al. Arch Biochem Biophys.* 1978 **185**:584. [PMID: 626512]
- [11] Trott O *et al. J Comput Chem.* 2010 **31**:455. [PMID: 19499576].
- [12] Morris GM *et al. J Comput Chem.* 2009 **30**:2785. [PMID: 19399780].

Edited by P Kanguane


Citation: Fathima *et al. Bioinformation* 17(1): 206-211 (2021)

License statement: This is an Open Access article which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. This is distributed under the terms of the Creative Commons Attribution License

Articles published in BIOINFORMATION are open for relevant post publication comments and criticisms, which will be published immediately linking to the original article for FREE of cost without open access charges. Comments should be concise, coherent and critical in less than 1000 words.

BIOINFORMATION

Discovery at the interface of physical and biological sciences



since 2005

BIOINFORMATION

Discovery at the interface of physical and biological sciences

indexed in

