Bioinformation 20(9): 1012-1016 (2024)

©Biomedical Informatics (2024)



Research Article



www.bioinformation.net Volume 20(9)

DOI: 10.6026/9732063002001012

Received September 1, 2024; Revised September 30, 2024; Accepted September 30, 2024, Published September 30, 2024

BIOINFORMATION

Discovery at the interface of physical and biological sciences

BIOINFORMATION 2022 Impact Factor (2023 release) is 1.9.

Declaration on Publication Ethics:

The author's state that they adhere with COPE guidelines on publishing ethics as described elsewhere at https://publicationethics.org/. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues connecting to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher in regard to this article.

Declaration on official E-mail:

The corresponding author declares that lifetime official e-mail from their institution is not available for all authors

License statement:

This is an Open Access article which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. This is distributed under the terms of the Creative Commons Attribution License

Comments from readers:

Articles published in BIOINFORMATION are open for relevant post publication comments and criticisms, which will be published immediately linking to the original article without open access charges. Comments should be concise, coherent and critical in less than 1000 words.

Disclaimer:

The views and opinions expressed are those of the author(s) and do not reflect the views or opinions of Bioinformation and (or) its publisher Biomedical Informatics. Biomedical Informatics remains neutral and allows authors to specify their address and affiliation details including territory where required. Bioinformation provides a platform for scholarly communication of data and information to create knowledge in the Biological/Biomedical domain.

> Edited by P Kangueane Citation: Kumari *et al.* Bioinformation 20(9): 1012-1016 (2024)

Comparison of CBCT examined root thickness and fracture resistance

Garima Kumari¹, Sibani Sarangi², Bandana Panda³, Vanka Aruna⁴, Jyoti Kasana⁵, Priya Sinha⁶ & Ritik Kashwani^{*,7}

¹Department of Conservative Dentistry and Endodontics, Manav Rachna Dental College, MRIIRS, Faridabad, India; ²Department of Periodontology, Hitech Dental College and Hospital, Bhubaneswar, India; ³Department of Conservative Dentistry and Endodontics, Kalinga Institute of Dental sciences, Bhubaneswar, India; ⁴Department of Periodontology, Bangalore Institute of Dental Sciences, Karnataka, India; ⁵North DMC Medical College and Hindu Rao Hospital, Delhi, India; ⁶Buddha Institute of Dental Sciences and Hospital, Magadh University, Patna, India; ⁷ Department of Oral Medicine and Maxillofacial Radiology, School of Dental Sciences, Sharda University, Greater Noida; *Corresponding author

Affiliation URL:

https://mriirs.edu.in/international-institute-of-research-and-studies/fds/ https://hi-techdentalbbsr.org/ https://kids.kiit.ac.in/

ISSN 0973-2063 (online) 0973-8894 (print)

Bioinformation 20(9): 1012-1016 (2024)

https://www.bids.edu/ https://hindurao.com/ https://bidsh.org/

Author contacts:

Garima Kumari - E-mail: cool.garima86@gmail.com Sibani Sarangi - E - mail: sibanisarangi94@gmail.com Bandana Panda -E - mail: iambandanapanda@gmail.com Vanka Aruna - E - mail: aruna.vankaz@gmail.com Jyoti Kasana - E - mail: kasanajt@gmail.com Priya Sinha - E - mail: dr.priyasinha02@gmail.com Ritik Kashwani - E - mail: docritikkashwani@yahoo.com

Abstract:

The correlation between remaining dentin thickness and fracture resistance in prepared teeth is of interest to dentists. A sample of 60 human teeth (mandibular premolars) was extracted and examined using cone beam computed tomography to assess residual dentin thickness before and after instrumentation. The gathered samples have been divided into three distinct categories, each with 20 samples. Hand files were used in Group 1, Protaper Next was used in Group 2, and a V taper was used in Group 3. They were examined for remaining dentin thickness after preparation with 3D CS Software and fracture resistance utilizing a Universal Testing Machine (UTM). Therefore, the objective of this research is to compare the amount of dentin removed by V Taper and Pro Taper Next to hand files using CBCT at the apical and coronal middle levels. Following that, these values will be correlated with fracture resistance values obtained from the Universal Testing Machine.

Keywords: CBCT, fracture resistance, Ni-Ti files, residual root thickness, universal testing machine.

Background:

Mechanical preparation of root canals using Ni-Ti rotary instruments causes vertical root fracture (VRF), which is a major problem associated with contemporary endodontic practice. Clinically, 10.9% to 31% of cases necessitate extraction either before or following endodontic therapy because of vertical root fracture [1, 2]. Among the multiple predisposing factors for VRF, the diameter of the prepared canal and the excessive taper of the rotary instruments were frequently noted as possible explanations for excessive dentin removal and root weakness; nevertheless, these remain debatable. All rotary systems used in dentistry have variable blade design, tip diameter, and configuration; however there is an important link between the Ni-Ti file system and dentinal micro fractures that lead to vertical root fracture [3, 4]. Bier et al. observed that root canal preparation with nickel titanium devices produced higher dentinal damage than by hand files [5]. Protaper Next with its off-set blade combined with the increasing and decreasing taper has shown to prevent taper lock thus resulting in fewer dentinal micro cracks. V Taper system with its variable taper design creates conservative coronal shape thus resulting in less dentinal removal and probably fewer dentinal micro cracks [6]. A handful of studies have been done on the dentin removal and fracture resistance of Pro Taper Next rotary instruments; however, no study has been reported to assess the fracture resistance of V taper rotary instruments. Numerous investigations verified the correlation between dentinal microcracks and vertical root fractures as a result of using various Ni-Ti files for instrumentation [7, 8]. Therefore, it is of interest to report the comparison of CBCT examined root thickness and fracture resistance.

Material and Methods:

This study was approved by the Institutional Research Ethical Committee at Manav Rachna Dental College (Approval ID ACAD/2017/492) and was conducted in the Department of Conservative Dentistry and Endodontics, Manav Rachna Dental College, Faridabad. It was approved by the internal ethical committee. We acquired orthodontically extracted human mandibular premolar teeth from patients ranging in age from 17 to 24. According to the rules and protocols set forth by the Occupational Safety and Health Administration, the extracted teeth were collected, stored, sterilized, and handled. To rule out the possibility of multiple canals, radiographs were taken in the Buccolingual and Mesiodistal angulations. The teeth were inspected for any cracks or anomalies that would need their exclusion using the Dental Operating Microscope (Global Aseries TM Microscope) at 10X magnification. After that, the teeth were washed and preserved for a week in 0.5% sodium hypochlorite solution. Following this, the teeth were decoronated to provide 13mm uniform root lengths.

Sample size calculation and distribution:

With the study's power established at 80.0% [(type II error = 0.20]] and 5% Type 1 error probability [α =0.05)], a sample size of 60 was selected. Following that, using the online randomization tool www.randomiser.org, the chosen teeth were numbered and categorized into three experimental groups of 20 each.

Mounting of teeth:

10 silicon molds, each containing six teeth (Affinis Putty, Coltene Whale dent), were utilized for positioning the teeth. Lead films were set on the right side of the mold in order to aid with the ISSN 0973-2063 (online) 0973-8894 (print)

Bioinformation 20(9): 1012-1016 (2024)

alignment of the teeth in the CBCT sections. A third person who was not participating in the study aligned the samples and allocated dental codes.

CBCT imaging:

Each mould was subjected to the CBCT scanner (Giano. Newtom Italy), using the image protocol for teeth, with the following exposure parameter: 10 Kv, 8mA, and 9 seconds. Each section was evaluated at 4 points (mesial, distal, buccal and lingual) in CS3D CBCT software.

Preparation of teeth:

Under the supervision of a senior faculty member with over twelve years of postgraduate teaching experience, a single operator performed the biomechanical preparation of the teeth. Six teeth were instrumented simultaneously to eliminate operator bias and fatigue. A No. 2 round carbide bur was utilized to prepare or modify the endodontic access cavity, and an Endo Access bur (DENTSPLY Endo Access Bur FG 1) was utilized to refine it. A DG 16 explorer was employed to locate the canals (Hu Friedy, IL USA). A # 10 K file was pushed 1 mm past the apical foramen and then removed to assess the canal's patency and calculate its working length. One millimetre less than the anatomic apex were the final working length.

Group 1(n=20): Step-back technique using stainless steel K-files:

Using a quarter turn pull, K-files (Mani Inc., Japan) prepared the canals up to #40 as the master apical file (MAF), after which they stepped back in increments of 1 mm for the subsequent three sizes of files (#45, #50, and #55). Each step back size file was accompanied by recapitulation utilizing the MAF at the working length.

Group 2: Pro Taper Next rotary files:

Using a torque-controlled endo motor (E connect S, Oricam), canals were prepared using the Protaper Next system in accordance with manufacturer guidelines up to X4 (40.06) at 300 rpm and 2 Ncm torque.

Group 3: V taper rotary files (SS White):

Making use of the torque-controlled endo motor (E connect S, Oricam), canals were prepared up to size (40.06). After every filling, the canals were flushed with 5 milliliters of 17% EDTA for duration of one minute, after being irrigated with 2 milliliters of normal saline and 1 milliliter of 2.5% NaOCL.

Evaluation of remaining dentin thickness:

Following canal preparation, the samples were placed back into the original mold, and CBCT scans were taken using a methodology akin to that of the original imaging. The minimal residual thickness (MRRT) was subsequently assessed. The residual root thickness (RRT) was subtracted from the initial root thickness (IRT) to determine the amount of dentin removal (DR). With the aid of the CS 3D imaging program, the data was examined.

Evaluation of fracture resistance:

Precision balances and vernier calipers have been employed to measure the weight and Buccolingual (BL) and Mesiodistal (MD) dimensions of the roots, respectively, and to fix the measurement errors. For block preparation, acrylic resin was utilized. Each root's coronal portion was exposed about 9 mm because of the apical root end being placed 4 mm vertically in the acrylic resin. The root was fractured via universal testing equipment having a cross head speed of 1 mm/min. A conical tip made from steel with a tip diameter of 0.5 mm was fixed to each specimen and placed parallel to its long axis with the canal orifice in center. The load necessary for fracture was recorded and expressed in Newton (N).



Figure 1: Comparison graph of MRRT

Results:

Statistical analysis:

One-way ANOVAs were primarily utilized for statistical analysis, and for multiple comparisons, the Post-hoc test was implemented afterward. The data provided was shown as mean +SD. The change in relative value with regard to time was assessed using a paired t-test. P-values below 0.05 are, at a 95% confidence level, considered to be significant. The analysis was performed employing SPSS 18.0, statistical software.

Minimum residual root thickness (MRRT):

When MRRT was compared within the group, the coronal third showed a statistically significant difference whereas the middle and apical third revealed no significant change. (P<0.001) Groups 1 and 2 demonstrated a significant difference in MRRT intergroup comparison, whereas groups 1 and 3 indicated no discernible distinction. (P < 0.001) The comparative analysis of minimal residual root thickness among all the groups has been demonstrated in (Figure 1) (Table 1).

Dentin removal (DR):

There was no statistically significant distinction between groups 1 and 3, nonetheless there was a substantial variance between the two when it related to dentin removal (DR) in the intergroup comparison (P<0.001). With group 2, dentin removal seemed more conspicuous than with group 1, which was followed by group 3 (Table 2).

```
Group 2 >Group 3 > Group 1
```

Bioinformation 20(9): 1012-1016 (2024)

©Biomedical Informatics (2024)

tuble 1. The mean of the mixer (minimum residual root discriciss) at american sections of single root manabular premotars in an ance groups (in min)								
		Ν	MEAN	STD DEVIATION	MINIMUM	MAXIMUM	F-VALUE	P-VALUE
MRRT 9mm from apex	GROUP 1 Hand File	20	1.304	0.953	0.82	5.3		
	GROUP 2 Protaper	20	0.6	0.158	0.35	0.98	7.749 0.001	
	GROUP 3 VTaper	20	0.945	0.168	0.68	1.2		
	TOTAL	60	0.95	0.627	0.35	5.3		
MRRT 6mm from apex	GROUP 1 Hand File	20	1.3	0.147	1	1.58		
	GROUP 2 Protaper	20	1.228	0.581	0.65	3.5	0.205	0.815
	GROUP 3 VTaper	20	1.273	0.169	1.02	1.78		
	TOTAL	60	1.267	0.354	0.65	3.5		
MRRT 3mm from apex	GROUP 1 Hand File	20	1.514	0.167	1.2	1.8		
	GROUP 2 Protaper	20	1.419	0.18	1	1.65	2.536	0.088
	GROUP 3 VTaper	20	1.514	0.105	1.35	1.72		
	TOTAL	60	1.482	0.158	1	1.8		

Table 1: The mean of the MRRT (Minimum residual root thickness) at different sections of single root mandibular premolars in all three groups (in mm)

Table 2: The mean of the DR (dentin removal) at different sections of single root mandibular premolars in all three groups (in mm)

		IN	wiean	Stu. Deviation	Winninum	Maximum	r-value	p-value
DR 9mm from apex	Group 1 Hand file	20	0.166	0.089	0.08	0.32	71.603	< 0.001
	Group 2 Pro Taper	20	0.668	0.197	0.32	0.98		
	Group 3 V taper	20	0.246	0.118	0.1	0.65		
	Total	60	0.36	0.262	0.08	0.98		
DR 6mm from apex	Group 1 Hand file	20	0.199	0.114	0.08	0.42	18.43	< 0.001
	Group 2 Pro Taper	20	0.428	0.2	0.2	0.85		
	Group 3 V taper	20	0.196	0.067	0.1	0.35		
	Total	60	0.274	0.174	0.08	0.85		
DR 3mm from apex	Group 1 Hand file	19	0.143	0.082	0.08	0.38	3.66	0.032
	Group 2 Pro Taper	20	0.234	0.141	0.1	0.5		
	Group 3 V taper	20	0.175	0.084	0.08	0.35		
	Total	59	0.185	0.111	0.08	0.5		

Table 3: Cross sectional diameters, multiplication of the BL-MD Diameters, weights, and facture loads of the roots

Groups	N	BL	MD	Multiplication of BL and MB	Weight (g)	Fracture load (N)
Hand file	20	7.24 + 0.44	5.16 + 0.48	37.38+3.92	0.45 ± 0.05	153.55+0.02
Pro Taper file	20	7.16+0.39	5.18+0.36	37.12+3.53	0.43+0.03	73.95+0.6
V Taper	20	7.26+0.42	5.15+0.38	37.37+3.65	0.44 + 0.05	207.20+0.7



Figure 2: Graph of FR result

Fracture resistance:

During assessment, each root examined in this study demonstrated a vertical fracture extending in the labiolingual direction. The three groups varied substantially, based on the results. Group 2 possessed the smallest degree of fracture resistance, followed by Group 1 and Group 3 showed the greatest amount of fracture resistance. (Table 3) (Figure 2)

Group 3 >Group 1 > group 2

Discussion:

One of the most critical iatrogenic components that affect a tooth's capability to endure a fracture is the thickness of the residual dentin following root canal treatment. Recent advances in non-surgical endodontic treatment techniques have begun to support rotary instrumentation considerably. Research indicates that, in comparison to hand files made of stainless steel, rotary instruments made out of Ni-Ti alloy have improved biomechanical preparation, enhancing the efficacy of root canal preparation [9-10]. Nevertheless, these instruments can either entirely or partially remove dentin from teeth while cleaning and shaping, weakening the tooth structure that remains and increasing the likelihood of breakage. While utilizing a highly tapered instrument for dentin removal, Zandbiglari et al. determined that the teeth appeared more susceptible to fracture compared to if one used hand instruments during preparation [3]. According to Lertchirakkarn et al. (2003), vertical root fracture wasn't caused solely by forces applied while lateral compaction [12, 13]. Thus, without obturating the prepared teeth, the study's main emphasis was on evaluating the dentin thickness which remained and the vertical root fracture. One of the newest technological advancements in radiology that can be used for research in the dentistry and medical fields is CBCT. With the capability to determine the remaining dentin thickness before as well as following instrumentation, this device provides Bioinformation 20(9): 1012-1016 (2024)

an advantageous and non-invasive technique which holds great promise for applications in endodontic research. Thus, the primary objective of the present research investigation sought to assess the residual root thickness following instrumentation using hand and different rotary files using CBCT analysis, and investigate whether or not this thickness has an association with the tooth's capability to endure fracture. Both fracture resistance [14-21] and residual dentin thickness [22-27] have been evaluated individually in quite a few of studies that have been published. Only a few individuals looked at the two metrics' links [28]. Therefore, without obturating the prepared teeth, the study's main objective was on evaluating the dentin thickness which remained and vertical root fracture. Additionally, only a limited number of studies [28] have provided an in-depth assessment of dentin removal at the apical, middle, and coronal aspects – aspects which the present research has also assessed.

Protaper Next (PTN) and V Taper rotary instruments were used to prepare the canals as opposed to the hand-instrumented group. There are researches [22, 23, 24] that analyze PTN, but to as far as we have knowledge, no published research reviews the fracture resistance of V Taper rotary files. The company's representatives claim that the V Taper files' unique design is expected to result in less dentin loss, strengthening the teeth's resistance to breakage. The dentin removal for Group PTN was significantly distinct from V Taper and traditional files (P<0.001), as determined by the data. Group 1 and Group 3 were not substantially different from each other, indicating that the amount of dentin removed with a V-taper was roughly comparable to that obtained with hand files (Table 2). This is consistent with the studies conducted by Akhlagi et al. (2010) [6]. The more prominent taper could have contributed to the more dentin loss observed in the coronal section of the canals, judging by the data. The fracture resistance of the three groups differed substantially, based on the results (Table 3). The groups having the lowest fracture resistance were Group 2 (Pro Taper Next), Group 1 (Hand file), and Group 3 (V Taper), which showed the highest fracture resistance. This suggested that there is a significant relationship between vertical root fracture and residual dentin thickness. The null hypothesis was thus disproved. To further corroborate these findings, bigger numbers of samples for comparable research are encouraged. One of the drawbacks of this research was that, considering it was carried out in vitro, care had to be taken when evaluating the results for clinical use.

Conclusion:

The study found a positive correlation between remaining dentin thickness and fracture resistance of teeth. CBCT

examination revealed that V Taper rotary file systems preserved residual root thickness more effectively than hand instrumentation, with V Taper showing the highest fracture resistance. The findings suggest that rotary systems with conservative designs, such as V Taper, better preserve tooth structure and enhance fracture resistance during endodontic procedures.

References:

- [1] Fuss Z et al. Int Endod J. 1999 32:283. [PMID: 10551119]
- [2] Sjogren U et al. J Endod. 1990 16:498. [PMID: 2084204]
- [3] Zandbiglari T *et al. Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006 **101**:126. [PMID: 16360617]
- [4] Adorno CG et al. J Endod. 2011 37:522. [PMID: 21419302]
- [5] Bier CA et al. J Endod. 2009 35:236.[PMID: 19166781]
- [6] Akhlaghi NM *et al. Int Endod J.* 2010 **43**:1029.[PMID: 20636352]
- [7] Ganesh A et al. J Conserv Dent. 2014 17:61.[PMID: 24554864]
- [8] Bhagat A et al. J Contemp Dent Pract. 2017 18:1065. [PMID: 29109323]
- [9] Shemesh H et al. Int Endod J. 2009 42:208.[PMID: 19228210]
- [10] Tan BT & Messer HH. J Endod. 2002 28:658.[PMID: 12236311]
- [11] Portenier I et al. Int Endod J. 1998 31:103. [PMID: 9868936]
- [12] Lertchirakarn V et al. J Endod. 1999 25:99. [PMID: 10204465]
- [13] Lindauer PA et al. J Endod. 1989 15:345. [PMID: 2637325]
- [14] Uyanik OM et al. J Endod. 2006 32:668. [PMID: 16793477]
- [15] Nagarthana PJ et al. J Indian Soc Pedod Prev Dent. 2006
 24:186.[PMID: 17183182]
- [16] Plotino G et al. Int Endod J. 2007 40:852. [PMID: 17697107]
- [17] Mahran AH & Mona M. J Endod. 2008 34:1219.[PMID: 18793924]
- [18] Vaudt J et al. Int Endod J. 2009 42:22. [PMID: 19125977]
- [19] Akhlaghi NM et al. J Conserv Dent. 2014 17:142.[PMID: 24778510]
- [20] Homayoon A *et al. Iran Endod J.* 2015 10:263.[PMID: 26525955]
- [21] Jain A et al. J Conserv Dent. 2016 19:440. [PMID: 27656063]
- [22] Sathorn C et al. J Endod. 2005 31:283.[PMID: 15793385]
- [23] Kazandag MK *et al. Int Endod J.* 2009 **42**:705.[PMID: 19467043]
- [24] Kim CH et al. J Endod. 2010 36:1195.[PMID: 20630298]
- [25] Abou El Nasr HM *et al. J Endod*. 2014 **40**:849.[PMID: 24862715]
- [26] De-Deus G et al. J Endod. 2015 41:676.[PMID: 25670245.]
- [27] Nur BG et al. Eur J dent. 2015 9:189.[PMID: 26038648]
- [28] Ibrahim AM et al. J Prosthet dent. 2016 115:290 [PMID: 26548871]