



www.bioinformation.net  
Volume 20(9)



Research Article

Received September 1, 2024; Revised September 30, 2024; Accepted September 30, 2024, Published September 30, 2024

DOI: 10.6026/973206300200990

BIOINFORMATION 2022 Impact Factor (2023 release) is 1.9.

**Declaration on Publication Ethics:**

The author's state that they adhere with COPE guidelines on publishing ethics as described elsewhere at <https://publicationethics.org/>. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues connecting to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher in regard to this article.

**Declaration on official E-mail:**

The corresponding author declares that lifetime official e-mail from their institution is not available for all authors

**License statement:**

This is an Open Access article which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. This is distributed under the terms of the Creative Commons Attribution License

**Comments from readers:**

Articles published in BIOINFORMATION are open for relevant post publication comments and criticisms, which will be published immediately linking to the original article without open access charges. Comments should be concise, coherent and critical in less than 1000 words.

**Disclaimer:**

The views and opinions expressed are those of the author(s) and do not reflect the views or opinions of Bioinformation and (or) its publisher Biomedical Informatics. Biomedical Informatics remains neutral and allows authors to specify their address and affiliation details including territory where required. Bioinformation provides a platform for scholarly communication of data and information to create knowledge in the Biological/Biomedical domain.

Edited by P Kanguane

Citation: Bhimsaria *et al.* Bioinformation 20(9): 990-992 (2024)

# An anatomico-morphometric analysis of proximal femur

Gaurangi Bhimsaria, T. Nagaeswari, T. Srimathi\* & Kalpana Ramachandran

Department of Anatomy, Sri Ramachandra Medical College and Research Institute, SRIHER (DU), Porur, Chennai 600 116, Tamil Nadu, India; \*Corresponding author

**Affiliation URL:**

<https://www.sriramachandra.edu.in/>

**Author contacts:**

Gaurangi Bhimsaria - E-mail: m0121168@sriher.edu.in, Phone-+91 7003870901

T Nagaeswari - E-mail:m0121214@sriher.edu.in, Phone-+91 80725 12592

T Srimathi - E-mail: drtsanatsrmc@gmail.com; srimathi.t@sriramachandra.edu.in, Phone: +9173584 41780

Kalpana Ramachandran - E-mail: hod.anatomy@sriramachandra.edu.in, Phone-+91 95661 37573

**Abstract:**  
The proximal femur is an important area of interest as it is involved in the articulation of the hip joint and it contributes to the mechanics of locomotion and the weight bearing capacity of the femur. Pathologies like osteoporosis and fractures are common in this site, and surgical interventions like hip arthroplasty and internal fixtures with implants rely on knowledge about the anatomy of the proximal femur to ensure effective treatment. Hence, a greater understanding of these variations will be helpful in constructing better fitted prosthesis. In this study, six morphometric parameters of the proximal femur were studied and measured in 100 dry adult femur bones - head diameter, transverse diameter of fovea, longitudinal diameter of fovea, diameter of neck, intertrochanteric line length, neck shaft angle and correlations were noted. The data obtained is specific for the southern Indian population and hopes to aid in production of better fitted hip prostheses for patients.

**Keywords:** Anthropometry; femur; femoral head; hip joint; hip prosthesis & morphometry

**Background:**

In the USA alone 5,44,000 hip arthroplasties are done on an average each year and in India the number is set to reach an all-time high by the year 2026. By the year 2050 the total number of hip fractures is projected to be double the number of cases confirmed in 2018. [1] With the rise in the number of surgeries involving the hip joint, a detailed analysis of the proximal femur and measurements of its parts will aid in creating finer adjusted and better suited prostheses in addition to implants which will in turn help in improving treatment outcomes. Various studies conducted in contemporary times have shown that these parameters of the femur are affected by a person's ethnicity, gender, lifestyle and environmental factors. Studies conducted in various parts of India have noted a difference in values due to influence of such factors [2, 3]. Therefore, it is of interest to observe the proximal end of femur and to report variations present proximal end of femurs of southern Indian population along with other correlations.

**Methodology:**

The materials used are Vernier Caliper, Goniometer and Dry femur bones. 100 dry adult femur bones are taken from Department of Anatomy in Sri Ramachandra Medical College. for the study, nature of which is observational and descriptive. Adult femurs which are intact, dried and non-pathological were included and femurs which have any arthritic deformity or gross damage were excluded in this study. Bones of the right and left side will be separated. The parameters measured are the diameter of head of femur (the linear distance measured between the upper and lower end of a femoral head in the cranio-caudal axis), transverse diameter of fovea (fullest extent of fovea capitis along the transverse axis), longitudinal diameter of fovea (fullest extent of fovea capitis along vertical axis), diameter of neck (the linear distance measured between the upper and lower end of the anatomical neck in the cranio-caudal axis), length of the intertrochanteric line (the total extent lengthwise of the intertrochanteric line) and the neck shaft angle with help of a vernier calliper with a least count of 0.01mm and goniometer with a least count of 1 degree. The values obtained will be recorded in Microsoft Excel. Then the values will be rounded up to two decimals and the mean along with standard deviation will be calculated. The data was tabulated and its statistical analysis was done using SPSS software. The findings of the six parameters measured are recorded in Table 2.

Table 1: Statistical summary of the proximal femur's morphometry

Parameter	N	Mean	Std Dev	Minimum	Maximum
Head Diameter	100	38.16	3	30	43
Transverse Diameter of Fovea	100	10.97	2.2	4	21
Longitudinal Diameter of Fovea	100	9.46	2.24	4	16
Diameter of Neck	100	28.62	3.33	21	36
Intertrochanteric Length	100	54.14	5.91	39	70
Neck-Shaft Angle	100	133.9	5.95	122	153

Table 2: Morphometric variables of right and left side femurs

Parameter	Group	N	Mean	Standard Deviation	S.E. Mean
Head Diameter	L	50	38.32	2.75	0.39
	R	50	38	3.25	0.46
Transverse Diameter of Fovea	L	50	11.61	1.92	0.27
	R	50	10.34	2.29	0.32
Longitudinal Diameter of Fovea	L	50	9.85	2.02	0.29
	R	50	9.07	2.39	0.34
Diameter of Neck	L	50	28.95	3.14	0.44
	R	50	28.29	3.51	0.5
Intertrochanteric Length	L	50	52.09	5.11	0.72
	R	50	56.19	6	0.85
Neck-Shaft Angle	L	50	132.4	5.36	0.76
	R	50	135.38	6.19	0.88

Table 3: Correlation between head diameter of a femur and longitudinal diameter of fovea

		Head Diameter	Longitudinal Diameter of Fovea
Head Diameter	Pearson Correlation	1	0.511
	Sig. (2-tailed)		0
	N	100	100
Longitudinal Diameter of Fovea	Pearson Correlation	0.511	1
	Sig. (2-tailed)	0	
	N	100	100

Table 4: Correlation between head diameter and neck diameter of proximal femur

		Head Diameter	Neck Diameter
Head Diameter	Pearson Correlation	1	0.735
	Sig. (2-tailed)		0
	N	100	100
Neck Diameter	Pearson Correlation	0.735	1
	Sig. (2-tailed)	0	
	N	100	100

Table 5: Comparison between different studies of proximal femur morphometry in Indian population

Parameter	Verma <i>et al.</i> [2]	Gupta <i>et al.</i> [3]	Lingamdenne PE <i>et al.</i> [4]	Present Study
Head Diameter (in mm)	42.32±4.11	41.59±3.25	42.3±5.40	38.16±3.00
Neck Diameter (in mm)	33.02±4.22	29.45±3.33	24.8±2.30	28.62±3.33
Neck Shaft Angle (in degrees)	128.90°±4.49°	119.08°±5.18°	119.44°±4.13°	133.89°±5.95°

**Results:**

The head diameter of the femur was found to be  $38.16 \pm 3.00$  mm. The transverse diameter of the fovea was found to be  $10.97 \pm 2.20$  mm whereas the longitudinal diameter was found to be  $9.46 \pm 2.24$  mm. The neck diameter was found to be  $28.62 \pm 3.33$  mm. Intertrochanteric length was  $54.14 \pm 5.91$  mm. The angle between neck and shaft was found to be  $133.89^\circ \pm 5.95^\circ$  (**Table 1**). A positive correlation was noted between head diameter and longitudinal diameter of fovea (**Table 3**). A significant positive correlation was also noted between the head diameter and neck diameter (**Table 4**).

**Discussion:**

Various methods have been used to measure the bony proximal femur's mentioned parameters—cadaveric morphometry, computed tomography, ultrasound and magnetic resonance imaging. The results obtained in this study are similar to other studies conducted in the Indian population (**Table 5**). Studies done in different population's revealed remarkable differences in these indices, showing that there is a need to customize prosthetic design that caters to every individual [2]. The fovea capitis is also an important anatomical structure in the proximal femur that transmits vessels supplying the femoral head. This ligament plays a role in cases where the head of femur undergoes avascular necrosis, which is a complication of hip fractures and dislocations. The mean transverse diameter of the fovea in our study was  $10.97 \pm 2.20$  mm. This value is comparable to those found by Gupta *et al.*, i.e.  $11.38 \pm 2.35$  mm. However, the longitudinal diameter of the fovea in our study,  $9.46 \pm 2.24$ , is much lesser than the value obtained by the same study  $15.94 \pm 3.37$  mm, suggesting a regional variation between northern and southern Indian populations. The computed tomography study done by Ceynowa *et al.* [5] in Poland found the transverse diameter to be  $12.94 \pm 2.61$  mm and the longitudinal diameter to be  $10.83 \pm 2.32$  mm, with the values being greater in men than in women. The intertrochanteric length was found to be  $54.14 \pm 5.91$  mm which is lesser than the value of  $60.31 \pm 7.33$  mm obtained in the dry bone study by Resmi George & Nithin K Raju. [6] The mean neck shaft angle obtained in this study is larger than the values obtained by Isaac B. *et al.*

[7] and Late SV & Keche H, [8] but were similar to the values obtained by Kamath SU *et al.* [9] and Haddad B *et al.* [10]

Interpretation of the values obtained in this study exhibit significant positive correlations between the longitudinal diameter of the fovea and the head diameter. Maximum positive correlation was seen between the neck diameter and the vertical head diameter.

**Conclusion:**

The morphometric analysis performed on the proximal femur provides valuable insights into anatomical variations and structural characteristics of the bone. The proximal femoral morphometry has notable differences among different groups of populations and such differences must be carefully analysed and taken into consideration while designing prostheses and in surgical interventions. The results obtained through this study will add to the existing database that will be helpful in designing implants, plates and prostheses for hip reconstructive surgeries that are catered to the Southern Indian population.

**References:**

- [1] Sing CW *et al.* *J Bone Miner Res.* 2023 **38**:1064 [PMID: 37118993].
- [2] Verma M *et al.* *J Clin Diagn Res.* 2017 **11**:AC01 [PMID: 28384844].
- [3] Gupta M *et al.* *Cureus.* 2022 **14**:e28780. [PMID: 36225441].
- [4] Lingamdenne PE & Marapaka P, *Indian J Clin Anat Physiol.* 2016 **3**:427 [DOI: 10.5958/2394-2126.2016.00097.9].
- [5] Ceynowa M *et al.* *Surg Radiol Anat.* 2019 **41**:101 [PMID: 30171297].
- [6] George R & Raju NK. *Asian Journal of Medical Sciences.* 2024 **15**:80. [doi.org/10.3126/ajms.v15i3.59822].
- [7] Isaac B *et al.* *Clin Anat.* 1997 **10**:318 [PMID: 9283729].
- [8] Late SV & Keche H. *Cureus.* 2022 **14**:e29188. [PMID: 36507110].
- [9] Kamath SU *et al.* *Malays Orthop J.* 2020 **14**:143. [PMID: 33403075].
- [10] Haddad B *et al.* *BMC Musculoskelet Disord.* 2022 **23**:1092. [PMID: 36514028].